

Functional Perspective of Feeder Organelle from Three-dimensional Ultrastructural Characteristics in *Cryptosporidium parvum*

著者	BOCHIMOTO Hiroki, KONDOH Daisuke, ISHIHARA Yo, Mohammad Hazzaz Bin KABIR, KATO Kentaro
journal or publication title	Journal of Integrated Field Science
volume	17
page range	1-4
year	2020-03
URL	http://hdl.handle.net/10097/00128787



Symposium mini review

Functional Perspective of Feeder Organelle from Three-dimensional Ultrastructural Characteristics in *Cryptosporidium parvum*

Hiroki BOCHIMOTO^{1,2}, Daisuke KONDOH³, Yo ISHIHARA⁴, Mohammad Hazzaz Bin KABIR²
and Kentaro KATO^{2,5}

¹Division of Aerospace Medicine, Department of Cell Physiology, The Jikei University School of Medicine, 3-25-8 Nishishimbashi, Minatoku, Tokyo 105-8461, Japan

²National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine, Nishi 2-13 Inada-cho, Obihiro, Hokkaido 080-8555, Japan

³Laboratory of Veterinary Anatomy, Obihiro University of Agriculture and Veterinary Medicine, Nishi 2-11 Inada-cho, Obihiro 080-8555, Japan

⁴Shonan Kamakura General Hospital, 1370-1 Okamoto, Kamakura, Kanagawa 247-8533, Japan

⁵Laboratory of Sustainable Animal Environment, Graduate School of Agricultural Science, Tohoku University, 232-3 Yomogida, Naruko-onsen, Osaki, Miyagi 989-6711, Japan

Keywords

Cryptosporidium, feeder organelle,
osmium maceration, SEM,
tubuloreticular formation

Corresponding Author

Hiroki BOCHIMOTO,
botimoto@jikei.ac.jp

Abstract

Cryptosporidium is a parasite causing extensive illness both in livestock and humans. Feeder organelle of *Cryptosporidium* is the multi-membranous structures localized on the parasite-host-cell interface that deprives nutrients from host cells. Although the feeder organelle has been summarized as a highly invaginated membranous structure, the three-dimensional fine structure remains unclear. Osmium-maceration procedure for scanning electron microscopy (OS-SEM) is one of the methods to enable visualization of the intracellular ultrastructure including depth direction information after removing soluble proteins. Recently, we investigated and assessed *C. parvum* possessed on the surface of ileal epithelial cells of mice by using transmission electron microscopy (TEM) and OS-SEM. By TEM observation, feeder organelles were recognized as aggregated structures of concentric-, vertically- and randomly-lined bars. Correspondingly, OS-SEM observation revealed the reticulated network of stacked flat bursiform, ring-shaped bursiform and reticulated tubular membranes. These findings of the three-dimensional ultrastructural characteristics of feeder organelle, which are more intricate than expected, may potentially reinforce the limited knowledge regarding the nature of this attachment interface and the functional mechanisms around extraction of nutrients.

Introduction

Cryptosporidium is a parasite responsible for highly contagious, and cryptosporidiosis in humans and animals (Ryan, Zahedi, & Paparini, 2016). Transmission of *Cryptosporidium* is most often by the fecal-oral route via contaminated water, food or fomites, or by direct ingestion of infected feces, resulting in the intestinal diarrhea (Yoshida *et al.*, 2007). Moreover, *C. parvum* is relatively resistant to chlorine at the levels used in potable water. Therefore, cryptosporidiosis infection occurs as a waterborne outbreak

with the potential to affect many people at once (Efstratiou, Ongerth, & Karanis, 2017; Mac Kenzie *et al.*, 1994; Yamamoto *et al.*, 2000). There is no effective medical treatment for either intestinal or biliary cryptosporidiosis, and in AIDS patients, the infection is rarely spontaneously cleared (Chen *et al.*, 1998; Theodos, Griffiths, D'Onfro, Fairfield, & Tzipori, 1998). In recent years, *Cryptosporidium* is a life-threatening opportunistic pathogen for children in developing countries containing Africa and Asia, who urgently need specific anti-cryptosporidial therapies (Elfadaly, Hassanain, Hassanain, Barakat, & Shaapan, 2018; HMG *et al.*, 2018; Kotloff *et al.*,

2013; Liu *et al.*, 2016; Platts-Mills *et al.*, 2015). Thus, the infection mechanism of *Cryptosporidium* should be elucidated for the development of effective precaution or treatment of cryptosporidiosis (Ryan *et al.*, 2016; Vanathy, Parija, Mandal, Hamide, & Krishnamurthy, 2017).

The infection mechanism of *cryptosporidium* to host cell

Life cycle of *Cryptosporidium* is completed in a single host, and infective sporozoites attach to and invade gastrointestinal epithelial cells to form a unique parasitophorous vacuole on top of the cells (O'Hara & Chen, 2011). The main site of contact between the maturing parasite and the host cell is an extensively folded membrane structure, called the feeder organelle (Zapata, Perkins, Riojas, Wu, & Le Blancq, 2002). The feeder organelle is peculiar and still largely uncharacterized structures (Sharling *et al.*, 2010). The feeder organelle is considered as the site at which nutrient uptake from the host cell cytoplasm occurs (Clode, Koh, & Thompson, 2015; Marcial & Madara, 1986; O'Donoghue, 1995). A *Cryptosporidium*-specific ATP-binding cassette, CpABC1, involved in transportation of various molecules (e.g.: metabolites and lipids) across membranes is localized to feeder organelles, indicating that these organelles are important for selective nutrient absorption (Perkins, Riojas, Wu, & Le Blancq, 1999; Zapata *et al.*, 2002). The ultrastructure of feeder organelles in *Cryptosporidium* has been examined using transmission electron microscopy (TEM) in murine models of infection and in livestock, and it has been described as "highly invaginated" (Al-Mathal & Alsalem, 2013; O'Hara & Chen, 2011; Pohlenz, Bemrick, Moon, & Cheville, 1978; Rosales, Arnedo, & Mascaró, 1998; Umemiya, Fukuda, Fujisaki, & Matsui, 2005; Valigurová, Hofmannová, Koudela, & Vávra, 2007). The feeder organelle structure was also seen in replicas as multiple membrane facets (Marcial & Madara, 1986). However, how they intake and transport molecules at an organelle level from the host cells remain uncertain.

Osmium-maceration for scanning electron microscopy to visualize 3D ultrastructures of feeder organelles

In contrast to TEM intracellular analysis, scanning electron microscopy (SEM) had been only used to analyze the three-dimensional (3D) surface structure of *Cryptosporidium* (Chen *et al.*, 1998; Pohlenz *et al.*, 1978; Umemiya *et al.*, 2005). The osmium-maceration procedure for SEM (OS-SEM) developed by Tanaka *et al.* in 1981 enables us to make a direct observation the intracellular endomembranous organelles by removing the cytoplasmic soluble proteins selectively from the cracked surface of the cells with diluted OsO₄ solution (Hanaki, Tanaka, & Kashima, 1985; Stowe, Fukudome, & Tanaka, 1986; Tanaka & Mitsushima, 1984; Tanaka & Naguro, 1981). OS-SEM is a very useful method that enables comparative observation with TEM. Recently, we introduced this excellent method, OS-SEM, to visualize the intracellular membranous organelles, especially feeder organelles in *Cryptosporidium* (Bochimoto, Kondoh, Ishihara, Kabir, & Kato, 2019). *Cryptosporidium* oocyst remained on the surface

of terminal ileum even after osmium maceration with 0.1% diluted OsO₄ (Fig. 1A). Moreover, we confirmed that the present OS-SEM clearly visualizes the intracellular structures of the parasite organelles including endoplasmic reticulum and some types of granules (Fig. 1B and C). By utilizing comparative observation of OS-SEM and TEM, we visualize the 3D ultrastructure of feeder organelle possessing three different components at the host-parasite interface: reticulated networks of stacked "dome-shaped" bursiform membranes (concentrically-spread type; cFO); networks of "ring-shaped" bursiform membranes concentrically surrounding "dome-shaped" membranes (vertically-lined type; vFO); reticulated "tube-shaped" membranes (randomly-scattered type; rFO, referred Fig. 1C) (Bochimoto *et al.*, 2019).

Discussion and future perspectives

Feeder organelles of *Cryptosporidium* have been considered as an invaginated membrane structure that functions to secure a large surface area (Fayer, 2008). However our visualization of the 3D ultrastructural characteristics of feeder organelle indicated that feeder organelles are more intricate and organized than was previously thought (Bochimoto *et al.*, 2019). Particularly, randomly tubule-reticular formation of rFO has an similarity to membrane traffic-associated organelles including trans-Golgi network and transitional ER (Hammond & Glick, 2000; Liendo & Joiner, 2000; Pelletier *et al.*, 2002). This finding makes us presume that the rFO has specific functions of membrane trafficking of absorbed nutrients (Fig. 1 D). In the future, the localization of the *Cryptosporidium*-specific transporter, like CpABC1, on each type of the feeder organelle components should be investigated to more clarify the mechanism of transport the nutrients. One of our concerns is that successful parasitism by *Cryptosporidium* needs intricate interactions between the host and the parasite (O'Hara & Chen, 2011). OS-SEM enables to visualize the intracellular 3D ultrastructure in not only *Cryptosporidium* but also the host cells (Fig. 1C), which may deal with this concern by comparative observation with TEM. Another concern is that the ultrastructural variation of feeder organelle components might depend on the life cycle stages of *Cryptosporidium*. Monoclonal antibodies, which recognize novel epitopes that could be recently used to define intracellular development (Wilke *et al.*, 2018), may resolve this concern. Regarding the structural nature of individual stages, cell attachment and invasion processes, the future lies in 3D imaging as indicated by Clode *et al.* (2015). Ultrahigh-resolution 3D imaging methods could be adopted to ATUMTome sectioning and serial imaging by SEM, block-face serial imaging by SEM, focused ion beam SEM (FIBSEM) and electron tomography by TEM. In addition of these methods, the "long ignored" OS-SEM method is a very powerful tool of analysis of intracellular ultrastructure associated with infection of *Cryptosporidium*.

Acknowledgements

This study was funded by grants-in-aid for Young Scientists, Scientific Research (B) and (C) and Scientific Research on Innovative Areas (3805) from the Ministry of




Fig. 1. Osmium-maceration scanning electron microscopic (OS-SEM) images of *Cryptosporidium parvum* attached to mouse ileal cells. [A] Surface of ileal epithelium. Arrowheads indicate outer surfaces images of parasites. [B] Higher magnified image of cross-section of ileal epithelial surface. An arrow indicates the oocyst. Intracellular structures of *Cryptosporidium* are colored red. Microvilli of host cells are colored blue. [C] OS-SEM images of feeder organelle. An arrow indicates the oocyst. Intracellular area and randomly-scattered type feeder organelle (rFO) of *Cryptosporidium* are colored red and green, respectively. Microvilli of host cells are colored blue. [D] Schematic illustration of three-dimensional ultrastructure of feeder organelle. cFO, concentrically-spread type feeder organelle; vFO, vertically-lined type feeder organelle. Bars=1 μ m.

Education, Culture, Science, Sports, and Technology (MEXT) of Japan; by the joint research program from the National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine; and by the Livestock Promotional Subsidy from the Japan Racing Association.

Dedication

We dedicate this paper to the memory of the late Professor Keiichi Tanaka M.D., Ph.D. (1926-2019), who pioneered the field of biological SEM.

References

- Al-Mathal, E. M. and A. A. Alsalem (2013) Pomegranate (*Punica granatum*) peel is effective in a murine model of experimental *Cryptosporidium parvum* ultrastructural studies of the ileum. *Experimental Parasitology*, 134(4): 482-494. DOI: 10.1016/j.exppara.2013.05.004
- Bochimoto, H., D. Kondoh, Y. Ishihara, M. H. B. Kabir and K. Kato (2019) Three-dimensional fine structure of feeder organelle in *Cryptosporidium parvum*. *Parasitology International*, 73: 101958. DOI: 10.1016/j.parint.2019.101958
- Chen, X. M., S. A. Levine, P. Tietz, E. Krueger, M. A. McNiven, D. M. Jefferson, M. M. Nicholas and N. F. LaRusso (1998) *Cryptosporidium parvum* is cytopathic for cultured human biliary epithelia via an apoptotic mechanism. *Hepatology (Baltimore, Md.)*, 28(4): 906-913. DOI: 10.1002/hep.510280402
- Clode, P. L., W. H. Koh and R. C. A. Thompson (2015) Life without a Host Cell: What is *Cryptosporidium*? *Trends in Parasitology*, 31(12): 614-624. DOI: 10.1016/j.pt.2015.08.005
- de Lucas H. M. G., N. S. Silveira, R. V. L. Q. dos Santos, L. E. B. Cordeiro, F. X. Barreto, M. de Lima Bueno, and P. H. M. de Oliveira. (2018) *Cryptosporidiosis: Clinical and Public Health. Journal of Epidemiology and Public Health Reviews*, 3(1): 1-6. DOI: 10.16966/2471-8211.160
- Efstratiou, A., J. E. Ongerth and P. Karanis (2017) Waterborne transmission of protozoan parasites: Review of worldwide outbreaks - An update 2011-2016. *Water Research*, 114(October): 14-22. DOI: 10.1016/j.watres.2017.01.036
- Elfadaly, H. A., N. A. Hassanain, M. A. Hassanain, A. M. Barakat and R. M. Shaapan (2018) Evaluation of primitive ground water supplies as a risk factor for the development of major waterborne zoonosis in Egyptian children living in rural areas. *Journal of Infection and Public Health*, 11(2): 203-208. DOI: 10.1016/j.jiph.2017.07.025
- Fayer, R. (2008) General Biology. In *Cryptosporidium and Cryptosporidiosis*, 2nd edition, R. Fayer and L. Xiao (Eds.), pp. 1-42, Boca Raton, Florida, CRC Press.
- Hammond, A. T. and B. S. Glick (2000) Dynamics of transitional endoplasmic reticulum sites in vertebrate cells. *Molecular Biology of the Cell*, 11(9): 3013-3030. DOI: 10.1091/mbc.11.9.3013
- Hanaki, M., K. Tanaka and Y. Kashima (1985) Scanning electron microscopic study on mitochondrial cristae in the rat adrenal cortex. *Journal of Electron Microscopy*, 34(4): 373-380. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/3837809>
- Kotloff, K. L., J. P. Nataro, W. C. Blackwelder, D. Nasrin, T. H. Farag, S. Panchalingam, Y. Wu, S. O. Sow, D. Sur, R. F. Breiman, A. S. Faruque, A. K. Zaidai, D. Saha, P. L. Alonso, B. Tamboura, D. Sanogo, U. Onwuchekwa, B. Manna, T. Ramamurthy, S. Kanungo, J. B. Ochieng, R. Omure, J. O. Onundo, A. Hossain, S. K. Das, S. Ahmed, S. Qureshi, F. Quadri, R. A. Adegbola, M. Antonio, M. J. Hossain, A. Akinsola, I. Mandomando, T. Nhampossa, S. Acácio, K. Biswas, C. E. O'Reilly, E. D. Mintz, L. Y. Berkeley, K. Muhsen, H. Sommerfelt, R. M. Robins-Browne and M. M. Levine (2013)

- Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *The Lancet (London, England)*, 382(9888): 209-222. DOI: 10.1016/S0140-6736(13)60844-2
- Liendo, A. and K. A. Joiner (2000) Toxoplasma gondii: conserved protein machinery in an unusual secretory pathway? *Microbes and Infection*, 2(2): 137-144. DOI: 10.1016/S1286-4579(00)00274-4
- Liu, L., S. Oza, D. Hogan, Y. Chu, J. Perin, J. Zhu, J. E. Lawn, S. Cousens, C. Mathers and R. E. Black (2016) Global, regional, and national causes of under-5 mortality in 2000-15: an updated systematic analysis with implications for the Sustainable Development Goals. *The Lancet (London, England)*, 388(10063): 3027-3035. DOI: 10.1016/S0140-6736(16)31593-8
- Mac Kenzie, W. R., N. J. Hoxie, M. E. Proctor, M. S. Gradus, K. A. Blair D. E. Peterson, J. J. Kazmierczak, D. G. Addiss, K. R. Fox, J. B. Rose and J. P. Davis (1994) A massive outbreak in Milwaukee of cryptosporidium infection transmitted through the public water supply. *The New England Journal of Medicine*, 331(3): 161-167. DOI: 10.1056/NEJM199407213310304
- Marcial, M. A. and J. L. Madara (1986) Cryptosporidium: cellular localization, structural analysis of absorptive cell-parasite membrane-membrane interactions in guinea pigs, and suggestion of protozoan transport by M cells. *Gastroenterology*, 90(3): 583-594. DOI: 10.1016/0016-5085(86)91112-1
- O'Donoghue, P. J. (1995) Cryptosporidium and cryptosporidiosis in man and animals. *International Journal for Parasitology*, 25(2): 139-195. DOI: 10.1016/0020-7519(94)E0059-V
- O'Hara, S. P. and X. M. Chen (2011) The cell biology of cryptosporidium infection. *Microbes and Infection*, 13(8-9): 721-730. DOI: 10.1016/j.micinf.2011.03.008
- Pelletier, L., C. A. Stern, M. Pypaert, D. Sheff, H. M. Ngô, N. Roper, C. Y. He, K. Hu, D. Toomre, I. Coppens, D. S. Roos, K. A. Joiner and G. Warren (2002) Golgi biogenesis in Toxoplasma gondii. *Nature*, 418(6897): 548-552. DOI: 10.1038/nature00946
- Perkins, M. E., Y. A. Riojas, T. W. Wu and S. M. Le Blancq (1999) CpABC, a Cryptosporidium parvum ATP-binding cassette protein at the host-parasite boundary in intracellular stages. *Proceedings of the National Academy of Sciences of the United States of America*, 96(10): 5734-5739. DOI: 10.1073/pnas.96.10.5734
- Platts-Mills, J. A., S. Babji, L. Bodhidatta, J. Gratz, R. Haque, A. Havt, B. J. J. McCormick, M. McGrath, M. P. Olortegui, A. Samie, S. Shakoor, D. Mondal, I. F. N. Lima, D. Hariraju, B. B. Rayamajhi, S. Qureshi, F. Kabir, P. P. Yori, B. Mufamadi, C. Amour, J. D. Carreon, S. A. Richard, D. Lang, P. Bessong, E. Mduma, T. Ahmed, A. A. M. Lima, C. J. Mason, A. K. M. Zaidi, Z. A. Bhutta, M. Kosek, R. L. Guerrant, M. Gottlieb, M. Miller, G. Kang, E. R. Houpt and MAL-ED Network Investigators. (2015) Pathogen-specific burdens of community diarrhoea in developing countries: a multisite birth cohort study (MAL-ED). *The Lancet Global Health*, 3(9): e564-75. DOI: 10.1016/S2214-109X(15)00151-5
- Pohlenz, J., W. J. Bemrick, H. W. Moon and N. F. Cheville (1978) Bovine cryptosporidiosis: a transmission and scanning electron microscopic study of some stages in the life cycle and of the host-parasite relationship. *Veterinary Pathology*, 15(3): 417-427. DOI: doi.org/10.1177/030098587801500318
- Rosales, M. J., T. Arnedo and C. Mascaró (1998) Ultrastructural details of Cryptosporidium parvum development in calf intestine. *Memorias Do Instituto Oswaldo Cruz*, 93(6): 847-850. DOI: 10.1590/s0074-02761998000600027
- Ryan, U., A. Zahedi and A. Paparini (2016) Cryptosporidium in humans and animals-a one health approach to prophylaxis. *Parasite Immunology*, 38(9): 535-547. DOI: 10.1111/pim.12350
- Sharling, L., X. Liu, D. R. Gollapalli, S. K. Maurya, L. Hedstrom and B. Striepen (2010) A screening pipeline for antiparasitic agents targeting cryptosporidium inosine monophosphate dehydrogenase. *PLoS Neglected Tropical Diseases*, 4(8): e794. DOI: 10.1371/journal.pntd.0000794
- Stowe, S., H. Fukudome and K. Tanaka (1986) Membrane turnover in crab photoreceptors studied by high-resolution scanning electron microscopy and by a new technique of thick-section transmission electron microscopy. *Cell and Tissue Research*, 245(1): 51-60. DOI: 10.1007/BF00218086
- Tanaka, K. and A. Mitsushima (1984) A preparation method for observing intracellular structures by scanning electron microscopy. *Journal of Microscopy*, 133(Pt 2): 213-222. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/6368836>
- Tanaka, K. and T. Naguro (1981) High resolution scanning electron microscopy of cell organelles by a new specimen preparation method. *Biomedical Research (Tokyo, Japan)*, 2, Suppl: 63-70. Retrieved from <https://scholar.google.co.jp/scholar?hl=ja&q=Tanaka%2C+K%2C+Naguro%2C+T%281981%29.+High+resolution+scanning+electron+microscopy+of+cell+organelles+by+a+new+specimen+preparation+method.+Biomed+Res+2%3A+Suppl%2C63-70.&btnG=&lr=#0>
- Theodos, C. M., J. K. Griffiths, J. D'Onfro, A. Fairfield and S. Tzipori (1998) Efficacy of nitazoxanide against Cryptosporidium parvum in cell culture and in animal models. *Antimicrobial Agents and Chemotherapy*, 42(8): 1959-1965. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9687390> <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC105716/pdf/ac001959.pdf>
- Umemiya, R., M. Fukuda, K. Fujisaki and T. Matsui (2005) Electron microscopic observation of the invasion process of Cryptosporidium parvum in severe combined immunodeficiency mice. *The Journal of Parasitology*, 91(5): 1034-1039. DOI: 10.1645/GE-508R.1
- Valigurová, A., L. Hofmannová, B. Koudela and J. Vávra (2007) An ultrastructural comparison of the attachment sites between Gregarina steini and Cryptosporidium muris. *The Journal of Eukaryotic Microbiology*, 54(6): 495-510. DOI: 10.1111/j.1550-7408.2007.00291.x
- Vanathy, K., S. C. Parija, J. Mandal, A. Hamide and S. Krishnamurthy (2017) Cryptosporidiosis: A mini review. *Tropical Parasitology*, 7(2): 72-80. DOI: 10.4103/tp.TP_25_17
- Wilke, G., S. Ravindran, L. Funkhouser-Jones, J. Barks, Q. Wang, K. L. VanDussen, T. S. Stappenbeck, T. B. Kuhlenschmidt, M. S. Kuhlenschmidt and L. D. Sibley (2018) Monoclonal Antibodies to Intracellular Stages of Cryptosporidium parvum Define Life Cycle Progression In Vitro. *MSphere*, 3(3): 1-14. DOI: 10.1128/msphere.00124-18
- Yamamoto, N., K. Urabe, M. Takaoka, K. Nakazawa, A. Gotoh, M. Haga, H. Fuchigami, I. Kimata and M. Iseki (2000) Outbreak of cryptosporidiosis after contamination of the public water supply in Saitama Prefecture, Japan, in 1996. *Kansenshogaku Zasshi. The Journal of the Japanese Association for Infectious Diseases*, 74(6): 518-526. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10916342>
- Yoshida, H., M. Matsuo, T. Miyoshi, K. Uchino, H. Nakaguchi, T. Fukumoto, Y. Teranaka and T. Tanaka (2007) An outbreak of cryptosporidiosis suspected to be related to contaminated food, October 2006, Sakai City, Japan. *Japanese Journal of Infectious Diseases*, 60(6): 405-407. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/18032847>
- Zapata, F., M. E. Perkins, Y. A. Riojas, T. W. Wu and S. M. Le Blancq (2002) The Cryptosporidium parvum ABC protein family. *Molecular and Biochemical Parasitology*, 120(1): 157-161. DOI: 10.1016/S0166-6851(01)00445-5